

Black Crappie and White Crappie (from NC WINS) ^[1]

Black Crappie and White Crappie

Pomoxis nigromaculatus and Pomoxis annularis

Written by Brad E. Hammers
updated by N. Corey Oakley
[North Carolina Wildlife Resources Commission](#) ^[2].



Classification White Crappie and Black Crappie

Class: Osteichthyes (bony fishes)
Order: Perciformes
Family: Centrarchidae

Average Size

White crappie range from 6.7 to 20.9 in. in total length; black crappie range from 5.1 to 19.3 in. Typical catches of either, range from 8 to 10 in. Crappies can weigh up to 3 or 4 lbs., but they typically reach a weight of 1/4 to 1/2 lb.

Food

Zooplankton, crustaceans, insects and fish. Young crappies feed primarily on zooplankton and insects. Adults feed primarily on fish, but also feed on insects.

Spawning


Begins in March and continues through May. Crappies build nests in shallow, calm water near vegetation. The nests are often built in groups. The males guard the nests until the eggs hatch and the young leave. Black crappie females produce between 11,000 and 188,000 eggs; white crappie females produce between 2,900 and 91,700 eggs. Eggs for both species hatch within 3 to 5 days.

Young

Crappie fry feed on yolk sac for 3 to 5 days, and then they leave the nest. Maturity is reached at an age of 2 to 3 years.

Life Expectancy

A maximum age of about 9 years.




Range and Distribution

Crappies are native to North Carolina. Both crappie species move to shallow water to mate in the spring, making them easier for anglers to catch. Some people think the crappie is the best-tasting fish to eat!

General Information

Papermouth, “sac-a-lait” (bag of milk) and speckled perch are just a few of the nicknames people have given the crappie. Two species found throughout North America are the black crappie (*Pomoxis nigromaculatus*) and the white crappie (*Pomoxis annularis*). They are members of the sunfish family, which also includes largemouth bass and bluegill. Crappies are two of the largest of the panfish species. Over the years, their sportiness on light tackle and their quality as “tablefare” have made them increasingly popular among freshwater anglers. Today, the black and white crappies are among the most fished-for species throughout North Carolina and much of the southeastern United States. About one-third of all sportfish harvested are crappies.



DescriptionDiagram of the Black and White crappie

The black crappie is distinguished by irregularly spaced black blotches on its silvery-green to yellowish sides. In the white crappie these blotches often form vertical bars. The black crappie has a dark green, olive-colored back and a thin, compressed body, which is somewhat deeper than that of white crappie. Its dorsal and anal fins are about the same shape and size and are colored with rows of greenish black spots. The black crappie has 7 or 8 spiny dorsal fin rays, while the white crappie only has 5 or 6. Because crossbreeding sometimes occurs between black and white crappies and water quality often affects fish coloration, counting spiny dorsal fin rays is the best method for distinguishing between the two species.

History and Status

Crappies are native to North Carolina and can be found in most waters throughout the state. They are quite hardy and tolerate a wide range of environmental conditions. In North Carolina, the black crappie flourishes in Piedmont reservoirs and Coastal Plain rivers. The white crappie flourishes in the cloudy, muddy reservoirs of the Yadkin River watershed.

Habitat and Habits

The black crappie is usually found in cool, clear ponds, reservoirs and slow-moving rivers, while white crappie prefer more turbid waters. Crappie tend to congregate around areas with abundant aquatic vegetation, brush piles or other types of cover.


Crappies have an extraordinarily high reproductive rate. The spawning season for both species in North Carolina lasts from about March through May, when water temperatures warm to the 60- to 68-degree range, and the white crappie spawns when the water temperatures reach the 58- to 64-degree range. Males select the nest sites and clear circular beds 8 to 15 in. in diameter. The beds are often located in water that is 3 to 8 ft. deep. Black crappie nests are frequently found in gravel or in soft, muddy river or lake bottoms. White crappie nests are often located near structures such as brush piles, stumps or rock outcrops. Each spawning black crappie female produces between 11,000 to 188,000 eggs,

while each white crappie female produces 2,900 to 91,700 eggs. Males guard the eggs, which hatch within 3 to 5 days. After the eggs hatch, males continue to guard the fry (newly hatched fish) for a few days until the young leave the nest.

Microscopic animals, called zooplankton, and insects are the preferred foods of these young-of-the-year black and white crappies. On this diet, they can grow up to 4 in. during the first 12 months. When they reach 6 to 8 in., young crappie eat mostly smaller fish and minnows.

Adult crappies prefer forage fish found in the open-water areas of reservoirs, particularly thread-fin and gizzard shad. They are, however, often willing to feed on anything that fits into their mouths, including their own young. When little or no prey fish are available, which is often the case during the winter and early spring, adults will shift back to feeding on insects.

The black crappie grows slower than white crappie, but because of its stockiness, a black crappie will weigh more than a white crappie of similar size. The rate of growth depends on habitat, food availability and crappie population size for a given body of water. Too many crappies and not enough food results in slow growth or stunting. The tremendous reproductive capability of crappie species often results in stunting, particularly in small ponds.



Crappie are among the most fished-for species in North Carolina.

Crappie species are most encountered by anglers. Fishing for crappies is very popular in North Carolina, and fishermen consider crappies to be excellent sportfish. They can be caught on a variety of fishing equipment including cane poles, spinning and bait-casting outfits, fly rods and ice-fishing gear. Anglers usually try to locate a school of crappies and fish for them using minnows, small jigs or a combination of the two. Care is required when setting the hook or when landing the fish because its soft papery mouth tears easily, allowing hooks to pull out. Crappie species are most active and ea



Biologists evaluate crappie populations in many reservoirs and rivers throughout NC. sily caught at night and in the early morning.

NCWRC Interaction: How You Can Help

Biologists with the Commission's Division of Inland Fisheries evaluate crappie populations in many reservoirs and rivers throughout North Carolina. Many Piedmont reservoirs contain excellent crappie populations. The B. Everett Jordan ^[4] Reservoir in Chatham County ^[5] contains high numbers of crappie greater than 10 in. This fishery is a great example of cooperation between Commission biologists and the public to create a production fishery for now and for the future. NCWRC biologists have established that a 10-in. minimum size limit and a 20-fish-per-person daily creel limit were

needed to protect mature, adult crappie in this reservoir. Without the cooperation of anglers complying with the new regulations, B. Everett Jordan Reservoir would not be the outstanding crappie fishery that it is today.

References:

Harlan, James R. and Everett B. Speaker. Iowa Fish and Fishing (Iowa State Conservation Commission, 1956).
Menhinick, Edward F. The Freshwater Fishes of North Carolina (North Carolina Wildlife Resources Commission, 1991).
Tomelleri, Joseph R. and Mark E. Eberle. Fishes of the Central United States (University Press of Kansas, 1990).
Sternberg, Dick. The Compleat Freshwater Fisherman (Cy DeCosse Incorporated, 1987).

Credits:

Illustration by J.T. Newman. Photos by North Carolina Wildlife Resources Commission.

Produced by the Division of Conservation Education, Cay Cross—Editor.



Profile courtesy of NC WILDLIFE INFORMATION NETWORK SHARE (NC WINS)

Subjects:

Wildlife ^[6]

Authors:

Hammers, Brad E. ^[7]

Oakley, N. Corey ^[8]

Origin - location:

Coastal Plain ^[9]

Mountains ^[10]

Piedmont ^[11]

From:

NC Wildlife Information Network Share (NC WINS). ^[12]

11 June 2010 | Hammers, Brad E.; Oakley, N. Corey

Source URL: <https://www.ncpedia.org/black-crappie-and-white-crappie-nc?page=0>

Links

[1] <https://www.ncpedia.org/black-crappie-and-white-crappie-nc> [2] <https://www.ncwildlife.org/> [3] <https://creativecommons.org/licenses/by-nd/4.0/> [4] <https://www.ncpedia.org/biography/jordan-benjamin-everett> [5] <https://www.ncpedia.org/geography/chatham> [6] <https://www.ncpedia.org/category/subjects/wildlife> [7] <https://www.ncpedia.org/category/authors/hammers-brad-e> [8] <https://www.ncpedia.org/category/authors/oakley-n-corey> [9] <https://www.ncpedia.org/category/origin-location/coastal-pl> [10] <https://www.ncpedia.org/category/origin-location/mountains> [11] <https://www.ncpedia.org/category/origin-location/piedmont> [12] <https://www.ncpedia.org/category/entry-source/nc-wildlife-i>