#### **Growing Living Cells lab**

Problem: What factors influence the growth of cells in sterile media?

#### Background:

This experiment models how different variables can affect cell growth in a bioreactor. Media provides the nutrients that cells need to grow under optimum conditions. In this inquiry, you will pick one independent variable in the growth of your cells to influence. What you do with your cells will be entirely up to you and your research partner. Everyone will use the same types of containers for media growth and the same techniques.

#### **Procedure:**

The class will be growing yogurt bacteria in skim milk media. Yogurt bacteria have been cultured for centuries around the world and originated from wild bacteria. Yogurt bacteria produce lactic acid which denatures and precipitates milk proteins. Lactic acid will be the "product" we are trying to make in our biomanufacturing process.

You will be growing your culture in tubes. Every tube will receive 20 mL of sterile milk media and be inoculated with 2 mL of the "mother" culture (yogurt bacteria).

You will have one control group that you will do nothing to and you must have at least 3 test groups that you will use to test your independent variable in different amounts.

You will be evaluating the growth of your bacteria in three ways.

- 1) You will measure the pH before you inoculate, after you inoculate, after 24 hours and after 48 hours.
- 2) You will note changes in the culture using qualitative observations after 24 and 48 hours.
- 3) You will complete a serial dilution of one of your cultures and evaluate the growth on agar plates.

#### **Hypothesis:**

If I change	<b>,</b>
then the bacteria will grow (faster/slower)	
because	
Independent Variable:	
Test Groups: Control, 1, 2, 3	
Things to keep constant (things I won't change between the groups):	

Procedure:						
Materials:						
4 culture tubes	"Mother" culture	Skim Milk media				
pH paper	Sterile 20 mL and 2 m	L measurement tools				
Sterile swabs						
		s number, and date. Label	1. "Control", 2. "Group 1," 3.			
• •	"Group 2," 4. "Group 3."  . Add 20 mL of sterile milk media to each tube.					
5. To group 2. I wi						
7						
8						
9. Cap and mix we	ell by inverting several	times.				
10. Using a differer	nt sterile swab for each	tube, dip into the culture	and place some of the culture			
solution onto p	H paper and record the	e starting pH. Using your n	nicroscope, record any			
observations of	the culture.					
11. Loosen the caps	s to allow air flow and	place your culture tubes in	a cup in the incubator.			
12. After 24 hours,	12. After 24 hours, repeat step 10 and 11.					
13. After 48 hours,	13. After 48 hours, repeat step 10 and 11 and complete the "Serial Dilution" procedure.					
14. Discard cultures	14. Discard cultures as directed by your teacher.					
15. Record bacteria	al counts from your ser	ial dilution.				
Data:						
Quantitative Observation	ons:					
Group	Starting pH	24 hours pH	48 hours pH			
Control						
1						
3	_					
<u>-</u>						
40 hours Doots siel Centre	for Crown					
48 hour Bacterial Count	. for Group	==				

### **Qualitative Observations:**

Group	р	Starting observations		24 hou	24 hours obs.		48 hours obs.	
Contr	rol							
1								
2								
3								
Line (	Graph	<u>ı</u> :		i	oH vs. Time			
	14							
	13							
	12							
	11							
	10 9	+						
	8							
Hd	7							
	6							
	4							
	5 4 3 2							
	1							
	0							
			Starting pH		24 hour			our pH

Group 3

Data Analysis:

What happened to the pH for each of the groups?

Was it the same for all of the groups? If not, how was it different?

How were the observations for each of the groups different?
Was there bacterial growth in the tube you chose to sample? How many bacteria were present?
<u>Conclusions:</u>
Were there any differences between the control and the test groups?
Did your independent variable (what you tested) affect the dependent variable (cell growth)? How do you know?
Was your hypothesis supported by the data?
What mistakes did you make?
What other variables do you think could affect cell growth?
What would you change if you had a chance to do this experiment again? What do you think would help the cells grow or keep them from growing more than what you did?
What have you learned about the challenges of growing cells in a bioreactor to make a biological chemical?